

ON FINITE PRODUCT OF CONVOLUTIONS AND CLASSIFICATIONS OF HYPERBOLIC AND ELLIPTIC EQUATIONS

A. KILIÇMAN AND H. ELTAYEB

ABSTRACT. In this paper we consider the linear second order partial differential equation with non-constant coefficients; then by using the double convolution product we produce new equations with polynomials coefficients and we classify the new equations. It is shown that the classifications of hyperbolic and elliptic new equations are similar to the original equations that is the classification is invariant after finite double convolutions product.

1. INTRODUCTION

There is no general method to solve a PDE and the behavior of the solutions depend essentially on the classification of PDE therefore the problem of classification of partial differential equations is very natural and well known. Some of second-order linear partial differential equations can be classified as Parabolic, Hyperbolic or Elliptic. The classification provides a guide to appropriate initial and boundary conditions, as well as to smoothness of the solutions. If a PDE has coefficients which are not constant, it is possible that it will not belong to any of these categories but rather be of mixed type. Recently, A. Kılıçman and H. Eltayeb in [4], introduced a new method of classification for partial differential equations with polynomial coefficients. In this study we extend the current classification to the finite product of double convolution with variable coefficients. First of all we give the following definition since it has a relation with the present study.

Definition 1. If $F_1(x, y)$ and $F_2(x, y)$ are integrable functions then the following integral

$$F_1(x, y) *^x F_2(x, y) = \int_0^x F_1(x - \theta, y) F_2(\theta, y) d\theta$$

is called a single convolution with respect to x provided that the integral exists.

Similarly, if $F_1(x, y)$ and $F_2(x, y)$ are integrable functions then the following integral

$$F_1(\theta_1, \theta_2) * * F_2(\theta_1, \theta_2) = \int_0^y \int_0^x F_1(x - \theta_1, y - \theta_2) F_2(\theta_1, \theta_2) d\theta_1 d\theta_2$$

is known as double convolution provided that the integral exists, see the details in [3].

Let us consider the linear second order partial differential equation with non-constant coefficients in the form of

$$a(x, y)u_{xx} + b(x, y)u_{xy} + c(x, y)u_{yy} + d(x, y)u_x + e(x, y)u_y + f(x, y)u = 0 \quad (1.1)$$

and almost linear equation in two variable

$$au_{xx} + bu_{xy} + cu_{yy} + F(x, y, u, u_x, u_y) = 0 \quad (1.2)$$

Date: January 16, 2009.

2000 Mathematics Subject Classification. Primary 35L05; Secondary 44A35.

Key words and phrases. Hyperbolic equation, Elliptic equation, Double convolution and Classification of PDE.

where a, b, c , are polynomials and defined by

$$a(x, y) = \sum_{\beta=1}^n \sum_{\alpha=1}^m x^\alpha y^\beta, \quad b(x, y) = \sum_{j=1}^n \sum_{i=1}^m x^\zeta y^\eta, \quad c(x, y) = \sum_{l=1}^n \sum_{k=1}^m x^k y^l$$

and $(a, b, c) \neq (0, 0, 0)$ and the expression $au_{xx} + 2bu_{xy} + cu_{yy}$ is called the principal part of Eq (1.2), since the principal part mainly determines the properties of solution. If we multiply the Eq (1.2) by polynomial with single convolution as $k(x) *^x$ where $k(x) = \sum_{i=1}^m x^i$

then Eq (1.2) becomes

$$k(x) *^x [a(x, y)u_{xx} + b(x, y)u_{xy} + c(x, y)u_{yy} + F(x, y, u, u_x, u_y)] = 0 \quad (1.3)$$

where the symbol $*^x$ indicates single convolution with respect to x , and we shall classify more general form Eq (1.3) instead of Eq (1.2) by considering the function

$$D(x, y) = (k(x, y) *^x b(x, y))^2 - (k(x, y) *^x a(x, y))(k(x, y) *^x c(x, y)) \quad (1.4)$$

From Eq (1.4), one can see that if D is positive then Eq (1.4) it called Hyperbolic, if D is negative then Eq (1.4) it called Elliptic, otherwise parabolic.

First of all, we compute the coefficients of Eq (1.3) as follow

$$A(x, y) = k(x) *^x a(x, y) = \sum_{i=1}^m x^i *^x \sum_{\beta=1}^n \sum_{\alpha=1}^m x^\alpha y^\beta$$

by using single convolution definition and integral by part we obtain

$$A(x, y) = \sum_{\beta=1}^n y^\beta \int_0^x \sum_{i=1}^m \sum_{\alpha=1}^m (x - \mu)^i \mu^\alpha d\mu \quad (1.5)$$

by computing the integral in the right hand side of Eq (1.5) we have

$$\sum_{\beta=1}^n y^\beta \int_0^x \sum_{i=1}^m \sum_{\alpha=1}^m (x - \mu)^i \mu^\alpha d\mu = \sum_{\beta=1}^n \sum_{i=1}^m \sum_{\alpha=1}^m \frac{i! x^{\alpha+i+1} y^\beta}{((\alpha+1)((\alpha+2)\dots(\alpha+i+1))}$$

thus, we obtain the first coefficient of Eq (1.3) in the form of

$$A_1(x, y) = \sum_{\beta=1}^n \sum_{i=1}^m \sum_{\alpha=1}^m \frac{i! x^{\alpha+i+1} y^\beta}{((\alpha+1)((\alpha+2)\dots(\alpha+i+1))}$$

similarly, for the coefficients of the second part in Eq (1.3) we have

$$B_1(x, y) = \sum_{j=1}^n \sum_{\zeta=1}^m \sum_{i=1}^m \frac{i! x^{\zeta+i+1} y^\eta}{((\zeta+1)((\zeta+2)\dots(\zeta+i+1))}$$

also the last coefficient of Eq (1.3) given by

$$C_1(x, y) = \sum_{l=1}^n \sum_{k=1}^m \sum_{i=1}^m \frac{i! x^{k+i+1} y^l}{((k+1)((k+2)\dots(k+i+1))}$$

then one can easily set up

$$D_1(x, y) = B_1^2 - A_1(x, y)C_1(x, y). \quad (1.6)$$

for example in particular we can have

$$x^3 *^x x^2 y^3 u_{xx} + x^3 *^x x^3 y^4 u_{xy} + x^3 *^x x^4 y^5 u_{yy} = f(x, y) *^x g(x, y). \quad (1.7)$$

The first coefficients of Eq (1.7) given by

$$A_1(x, y) = x^3 *^x x^2 y^3 = y^5 \int_0^x (x - \theta)^3 \theta^2 d\theta = \frac{1}{60} y^3 x^6. \quad (1.8)$$

Similarly, the second coefficient given by

$$B_1(x, y) = x^3 *^x x^3 y^4 = \frac{1}{140} y^4 x^7. \quad (1.9)$$

By the same way we get the last coefficients of Eq (1.7)

$$C_1(x, y) = x^3 *^x x^4 y^5 = \frac{1}{280} y^5 x^8. \quad (1.10)$$

By using Eqs (1.6), (1.8), (1.9) and (1.10) we obtain

$$D_1(x, y) = -\frac{1}{117600} y^8 x^{14} \quad (1.11)$$

We can easy see from Eq(1.11) that Eq (1.7) is an elliptic equation for all (x_0, y) .

In the same way, if we multiply the Eq (1.2) by polynomial with single convolution as $h(y)*^y$ where $h(y) = \sum_{j=1}^n y^j$ then Eq (1.2) becomes

$$h(y) *^y [a(x, y)u_{xx} + b(x, y)u_{xy} + c(x, y)u_{yy} + F(x, y, u, u_x, u_y)] = 0 \quad (1.12)$$

where the symbol $*^y$ indicates single convolution with respect to y , and we shall classify Eq (1.12) as. First of all, let us compute the coefficients of Eq (1.12) by using definition of single convolution with respect to y and integral by part we obtain the first coefficient of Eq (1.12) as follow

$$A_2(x, y) = h(y) *^y a(x, y) = \sum_{\beta=1}^n \sum_{j=1}^n \sum_{\alpha=1}^m \frac{j! x^\alpha y^{\beta+j+1}}{((\beta+1)((\beta+2)\dots(\beta+j+1)))},$$

and the second coefficients of Eq(1.12) given by

$$B_2(x, y) = h(y) *^y b(x, y) = \sum_{j=1}^n \sum_{\eta=1}^n \sum_{i=1}^m \frac{j! x^\eta y^{\eta+j+1}}{((\eta+1)((\eta+2)\dots(\eta+i+1)))}.$$

Similarly, the last coefficients of Eq(1.12) given by

$$C_2(x, y) = h(y) *^y c(x, y) = \sum_{j=1}^n \sum_{l=1}^n \sum_{i=1}^m \frac{j! x^l y^{l+j+1}}{((l+1)((l+2)\dots(l+i+1)))}$$

In particular, let us classify the following example

$$y^7 *^y x^2 y^3 u_{xx} + y^7 *^y x^3 y^4 u_{xy} + y^7 *^y x^4 y^5 u_{yy} = f(x, y) *^y g(x, y). \quad (1.13)$$

the symbol $*^y$ means single convolution with respect to y . We follow the same technique that used above, then the first coefficient of Eq (1.13) given by

$$A_2(x, y) = y^7 *^y x^2 y^3 = \frac{1}{1320} x^2 y^{11}, \quad (1.14)$$

the second coefficient of (1.13) given by

$$B_2(x, y) = y^7 *^y x^3 y^4 = \frac{1}{3860} x^3 y^{12} \quad (1.15)$$

and the last coefficient given by

$$C_2(x, y) = y^7 *^y x^4 y^5 = \frac{1}{10296} x^4 y^{13}. \quad (1.16)$$

On using Eqs (1.18), (1.14), (1.15) and (1.16) we have

$$D_2(x, y) = -\frac{1}{101930400} x^6 y^{24} \quad (1.17)$$

We can easily see from Eq(1.17) that Eq (1.13) is an elliptic equation for all (x, y_0) .

Now, let us extend the above results form single convolution to double convolution as:

If we multiply the Eq (1.2) by polynomial with double convolution as $k(x, y) **$ where

$$k(x, y) = \sum_{j=1}^n \sum_{i=1}^m x^i y^j \text{ then Eq (1.2) becomes}$$

$$k(x, y) ** [a(x, y)u_{xx} + b(x, y)u_{xy} + c(x, y)u_{yy} + F(x, y, u, u_x, u_y)] = 0$$

where the symbol $**$ indicates double convolution and we shall classify more general form Eq (1.12) instead of Eq (1.2) by considering the function

$$D(x, y) = (k(x, y) ** b(x, y))^2 - (k(x, y) ** a(x, y)) (k(x, y) ** c(x, y)) \quad (1.18)$$

First of all, we compute the coefficients of Eq (1.18) as follow

$$A(x, y) = k(x, y) ** a(x, y) = \sum_{j=1}^n \sum_{i=1}^m x^i y^j ** \sum_{\beta=1}^n \sum_{\alpha=1}^m x^\alpha y^\beta$$

by using double convolution definition and integral by part we obtain

$$A(x, y) = \int_0^y \sum_{j=1}^n \sum_{\beta=1}^n (y - \nu)^j \nu^\beta d\nu \times \int_0^x \sum_{i=1}^m \sum_{\alpha=1}^m (x - \mu)^i \mu^\alpha d\mu \quad (1.19)$$

the first integral in the right hand side of Eq (1.19) can be obtain as

$$\int_0^y \sum_{j=1}^n \sum_{\beta=1}^n (y - \nu)^j \nu^\beta d\nu = \sum_{j=1}^n \sum_{\beta=1}^n \frac{j! y^{\beta+j+1}}{(\beta+1)((\beta+2)\dots(\beta+j+1))} \quad (1.20)$$

in a similar way the second integral in the right hand side of Eq (1.19) is given by

$$\int_0^x \sum_{i=1}^m \sum_{\alpha=1}^m (x - \mu)^i \mu^\alpha d\mu = \sum_{i=1}^m \sum_{\alpha=1}^m \frac{i! x^{\alpha+i+1}}{(\alpha+1)((\alpha+2)\dots(\alpha+i+1))} \quad (1.21)$$

thus from Eqs (1.20) and (1.21) we obtain the first coefficient of Eq (1.18) in the form of

$$A(x, y) = \sum_{j=1}^n \sum_{\beta=1}^n \sum_{i=1}^m \sum_{\alpha=1}^m \frac{i! j! x^{\alpha+i+1} y^{\beta+j+1}}{((\alpha+1)((\alpha+2)\dots(\alpha+i+1)) ((\beta+1)((\beta+2)\dots(\beta+j+1)))}$$

similarly, for the coefficients of the second part in Eq (1.18) we have

$$B(x, y) = \sum_{j=1}^n \sum_{l=1}^n \sum_{k=1}^m \sum_{i=1}^m \frac{i! j! x^{k+i+1} y^{l+j+1}}{((k+1)((k+2)\dots(k+i+1)) ((l+1)((l+2)\dots(l+j+1)))}$$

and similarly, the last coefficient of Eq (1.18) given by

$$C(x, y) = \sum_{j=1}^n \sum_{l=1}^n \sum_{k=1}^m \sum_{i=1}^m \frac{i! j! x^{\zeta+i+1} y^{\eta+j+1}}{((k+1)((k+2)\dots(k+i+1)) ((l+1)((l+2)\dots(l+j+1)))}$$

then one can easily set up

$$D(x, y) = B^2 - A(x, y)C(x, y). \quad (1.22)$$

From Eq (1.22), one can see that if D is positive then Eq (1.22) it called Hyperbolic, if D is negative then Eq (1.22) it called Elliptic, otherwise parabolic.

Generalized the Classification of Hyperbolic and Elliptic equations

In the this section we are going to generalized the classification of hyperbaric and elliptic equations as follow. Firstly, we consider the one dimensional wave equation with polynomial as follows

$$a(x, t)u_{tt} - c(x, t)u_{xx} = f_1(x, t) * f_2(x, t) \quad (1.23)$$

where a, b are polynomials defined by $a(x, t) = \sum_{\beta=1}^n \sum_{\alpha=1}^m x^\alpha t^\beta$ and $c(x, t) = \sum_{l=1}^n \sum_{k=1}^m x^\mu t^\nu$, now,

we generalize Eq (1.23) by using finite double convolutional product as follows

$$\prod_{r=1}^l p_r(x, t) * (a(x, t)u_{tt} - c(x, t)u_{xx}) = f_1(x, t) * f_2(x, t) \quad (1.24)$$

where $r = 1, 2, \dots, l$, and

$$\prod_{r=1}^l (p_r(x, t)) = p_1(x, t) * p_2(x, t) * p_3(x, t) * \dots * p_l(x, t)$$

where $p_1, p_2, p_3 \dots p_l$ polynomials defined by

$$p_1(x, t) = \sum_{j_1=1}^n \sum_{i_1=1}^m x^{i_1} t^{j_1}, \quad p_2(x, t) = \sum_{j_2=1}^n \sum_{i_2=1}^m x^{i_2} t^{j_2}, \dots, \quad p_l(x, t) = \sum_{j_l=1}^n \sum_{i_l=1}^m x^{i_l} t^{j_l} \quad (1.25)$$

To compute the first coefficients of Eq (1.24), we start by $l = 1$, we obtain the coefficient in the form

$$A_1(x, t) = p_1(x, t) * a(x, t) = \sum_{j_1=1}^n \sum_{i_1=1}^m x^{i_1} t^{j_1} * \sum_{\beta=1}^n \sum_{\alpha=1}^m x^\alpha t^\beta$$

by using double convolution definition and integral by part we obtain

$$A_1(x, t) = \int_0^t \sum_{j_1=1}^n \sum_{\beta=1}^m (t - \zeta)^{j_1} \zeta^\beta d\zeta \times \int_0^x \sum_{i_1=1}^m \sum_{\alpha=1}^m (x - \eta)^{i_1} \eta^\alpha d\eta \quad (1.26)$$

the first integral in the right hand side of Eq (1.26) can be obtain as

$$\int_0^t \sum_{j_1=1}^n \sum_{\beta=1}^m (t - \zeta)^{j_1} \zeta^\beta d\zeta = \sum_{j_1=1}^n \sum_{\beta=1}^m \frac{j_1! t^{\beta+j_1+1}}{(\beta+1)((\beta+2)\dots(\beta+j_1+1))} \quad (1.27)$$

in a similar way the second integral in the right hand side of Eq (1.26) is given by

$$\int_0^x \sum_{i_1=1}^m \sum_{\alpha=1}^m (x - \eta)^{i_1} \eta^\alpha d\eta = \sum_{i_1=1}^m \sum_{\alpha=1}^m \frac{i_1! x^{\alpha+i_1+1}}{(\alpha+1)((\alpha+2)\dots(\alpha+i_1+1))} \quad (1.28)$$

thus from Eqs (1.27) and (1.28) we obtain

$$A_1(x, t) = \sum_{j_1=1}^n \sum_{\beta=1}^m \sum_{i_1=1}^m \sum_{\alpha=1}^m x^{\alpha+i_1+1} t^{\beta+j_1+1},$$

where

$$Q = \frac{i_1! j_1!}{((\alpha+1)((\alpha+2)\dots(\alpha+i_1+1))((\beta+1)((\beta+2)\dots(\beta+j_1+1)))}$$

similarly, if we let $l = 2$, we have

$$A_2(x, t) = A_1(x, t) * p_1(x, t) = \sum_{j_1=1}^n \sum_{\beta=1}^m \sum_{i_1=1}^m \sum_{\alpha=1}^m Q x^{\alpha+i_1+1} t^{\beta+j_1+1} * \sum_{j_2=1}^n \sum_{i_2=1}^m x^{i_2} t^{j_2}$$

similarly, as above we obtain the coefficient in case $l = 2$, as follows

$$A_2(x, t) = \sum_{j_1=1}^n \sum_{j_2=1}^n \sum_{\beta=1}^n \sum_{i_1=1}^m \sum_{i_2=1}^m \sum_{\alpha=1}^m \frac{K}{GL} x^{\alpha+i_1+i_2+1} t^{\beta+j_1+j_2+1}$$

where

$$K = i_1!i_2!j_1!j_2!, \quad G = ((\alpha+1)(\alpha+2)\dots(\alpha+i_1+1))((i_1+\alpha+2)(i_1+\alpha+3)\dots(i_1+\alpha+1+i_2))$$

and $L = ((\beta+1)(\beta+2)\dots(\beta+j_1+1))(((j_1+\beta+2))(j_1+\beta+3))\dots((j_1+\beta+1+j_2)))$.

By the same way we compute the coefficient in case $l = 3$

$$A_3(x, t) = A_2(x, t) ** p_3(x, t) = \sum_{j_1=1}^n \sum_{j_2=1}^n \sum_{\beta=1}^n \sum_{i_1=1}^m \sum_{i_2=1}^m \sum_{\alpha=1}^m \frac{K}{GL} x^{\alpha+i_1+i_2+1} t^{\beta+j_1+j_2+1} ** \sum_{j_3=1}^n \sum_{i_3=1}^m x^{i_3} t^{j_3}$$

by using double convolution definition and integral by part we get

$$A_3(x, t) = \sum_{j_1=1}^n \sum_{j_2=1}^n \sum_{j_3=1}^n \sum_{\beta=1}^n \sum_{i_1=1}^m \sum_{i_2=1}^m \sum_{i_3=1}^m \sum_{\alpha=1}^m \frac{B}{MN} x^{\alpha+i_1+i_2+i_3+1} t^{\beta+j_1+j_2+j_3+1},$$

where

$$B = i_1!i_2!i_3!j_1!j_2!j_3!,$$

$$M = ((\alpha+1)(\alpha+2)\dots(\alpha+i_1+1))((i_1+\alpha+2)(i_1+\alpha+3)\dots(i_1+\alpha+1+i_2)) \\ \times (((i_1+i_2+\alpha+3)(i_1+i_2+\alpha+4)\dots(i_1+i_2+\alpha+1+i_3)))$$

and

$$N = ((\beta+1)(\beta+2)\dots(\beta+j_1+1))(((j_1+\beta+2))(j_1+\beta+3))\dots((j_1+\beta+1+j_2))) \\ \times ((j_1+j_2+\beta+3))\dots((j_1+j_2+\beta+1+j_3))$$

In general form the first coefficients of Eq (1.24) given by

$$A(x, t) = \prod_{r=1}^l p_r(x, t) ** (a(x, t)) \\ = \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_l=1}^n \sum_{i_l=1}^m \left(\sum_{\beta=1}^n \sum_{\alpha=1}^m \frac{\Phi\Psi}{\Theta\Xi} x^{i_1+i_2+\dots+i_l+\alpha+1} t^{j_1+j_2+\dots+j_l+\beta+1} \right) \quad (1.29)$$

where

$$\Phi = i_1!i_2!\dots i_l!, \quad \Psi = j_1!j_2!\dots j_l!, \\ \Theta = ((\alpha+1)(\alpha+2)\dots(\alpha+i_1+1))((i_1+\alpha+2)(i_1+\alpha+3)\dots(i_1+\alpha+1+i_2)) \dots \\ \dots ((i_1+\dots+i_{l-1}+\alpha+1)\dots(i_1+\dots+i_l+\alpha+1))$$

and

$$\Xi = ((\beta+1)(\beta+2)\dots(\beta+j_1+1))(((j_1+\beta+2))(j_1+\beta+3))\dots((j_1+\beta+1+j_2))) \dots \\ \dots ((j_1+j_2+\dots+j_{l-1}+\beta+1)\dots(j_1+j_2+\dots+j_l+\beta+1))$$

Similarly, the second coefficients of Eq (1.24) given by

$$C(x, t) = \prod_{r=1}^l p_r(x, t) ** (c(x, t)) \\ = \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_l=1}^n \sum_{i_l=1}^m \left(\sum_{\mu=1}^n \sum_{\nu=1}^m \frac{F\Upsilon}{\Delta\Lambda} x^{i_1+i_2+\dots+i_l+\mu+1} t^{j_1+j_2+\dots+j_l+\nu+1} \right) \quad (1.30)$$

where

$$\begin{aligned} F &= i_1!i_2!\dots i_l!, \quad \Upsilon = j_1!j_2!\dots j_l!, \\ \Delta &= ((\mu+1)(\mu+2)\dots(\mu+i_1+1))((\mu+i_1+2)(\mu+i_1+3)\dots(\mu+i_1+i_2+1))\dots \\ &\quad \dots((i_1+i_2+\dots+i_{l-1}+\mu+1)\dots(i_1+i_2+\dots+i_l+\mu+1)) \end{aligned}$$

and

$$\begin{aligned} \Lambda &= ((\nu+1)(\nu+2)\dots(\nu+j_1+1))((\nu+j_1+2)(\nu+j_1+3)\dots(\nu+j_1+j_2+1))\dots \\ &\quad \dots((j_1+j_2+\dots+j_{l-1}+\nu+1)\dots(j_1+j_2+\dots+j_l+\nu+1)) \end{aligned}$$

Now, we going back to Eq(1.22)

$$D(x, t) = -A(x, t)C(x, t).$$

Then we obtain $D(x, t)$ as follow

$$\begin{aligned} D(x, t) &= \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_l=1}^n \sum_{i_l=1}^m \left(\sum_{\beta=1}^n \sum_{\alpha=1}^m \frac{\Phi\Psi}{\Theta\Xi} x^{i_1+i_2+\dots+i_n+\alpha+1} t^{j_1+j_2+\dots+j_m+\beta+1} \right) \times \\ &\quad \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_l=1}^n \sum_{i_l=1}^m \left(\sum_{\mu=1}^n \sum_{\nu=1}^m \frac{F\Upsilon}{\Delta\Lambda} x^{i_1+i_2+\dots+i_n+\mu+1} t^{j_1+j_2+\dots+j_m+\nu+1} \right) \quad (1.31) \end{aligned}$$

We assume that all the coefficients $A(x, t)$, $C(x, t)$ are convergent. Now, we can consider some particular cases:

Case(1): If $i_1+i_2+\dots+i_l+j_1+j_2+\dots+j_l, \alpha+\beta$ and $\mu+\nu$ are odd, we classify Eq (1.24) by using Eq (1.31), since $\alpha+\beta$ and $\mu+\nu$ are odd, then the power of x, t in Eq (1.31) is even, thus we see that $D > 0$ for all $(x_0, t_0) \in \mathbb{R}^2$, Eq (1.24) Hyperbolic equation. Then for example if we consider a particular non-constant coefficient wave equation in one dimension in the form

$$x^5 t^2 ** (x^4 t^3 ** x^2 t^7) u_{tt} - x^5 t^2 ** (x^4 t^3 ** x^6 t^5) u_{xx} = f(x, t) ** g(x, t) \quad (1.32)$$

We follow the similar technique that was introduced by A. Kılıçman and H. Eltayeb, see [4], then in particular we compute the coefficients of Eq(1.32) by using MAPLE 11 as

$$A = x^5 t^2 ** x^4 t^3 ** x^2 t^7 = \frac{1}{1558311955200} x^{13} t^{14},$$

where

$$\begin{aligned} \Phi &= i_1!i_2! = 4!5! \\ \Psi &= j_1!j_2! = 3!2! \\ \Theta &= ((\alpha+1)(\alpha+2)\dots(\alpha+i_1+1))((i_1+\alpha+2)(i_1+\alpha+3)\dots(i_1+\alpha+2+i_2)) \end{aligned}$$

and $\Xi = ((\beta+1)(\beta+2)\dots(\beta+j_1+1))((j_1+\beta+2)(j_1+\beta+3)\dots(j_1+\beta+2+j_2))$, where $i_1 = 4, \alpha = 2, i_2 = 5, j_1 = 3, \beta = 7$ and $j_2 = 2$, thus

$$\frac{\Phi\Psi}{\Theta\Xi} = \frac{4!5!3!2!}{(3.4.5.6.7)(8.9.10.11.12.13)(8.9.10.11)(12.13.14)} = \frac{1}{1558311955200}$$

$$C = -x^5 t^2 ** x^4 t^3 ** x^6 t^5 = -\frac{1}{57058191590400} x^{17} t^{12},$$

similarly, the factor of second coefficients is given by

$$\frac{F\Upsilon}{\Delta\Lambda} = \frac{1}{57058191590400},$$

then

$$D = -AC = \frac{1}{88914462097412421550080000} x^{30} t^{26} \quad (1.33)$$

We can easily see from Eq(1.33) that Eq(1.32) is hyperbolic for all $(x_0, t_0) \in \mathbb{R}^2$.

Case(2): If $i_1 + i_2 + \dots + i_l + j_1 + j_2 + \dots + j_l, \alpha + \beta$ and $\mu + \nu$ are even in Eq (1.24), and assume $b = 0$, then the power of x, t in equation Eq (1.31) is even thus it follows that for all points (x_0, t_0) in the domain $\mathbb{R}^2$ Eq (1.24) is Hyperbolic.

Now, if we consider another particular non-constant coefficient wave equation in one dimension in the form of

$$xt^9 ** (x^5 t^3 ** x^3 t^7) u_{tt} - xt^9 ** (x^5 t^3 ** x^7 t^5) u_{xx} = f(x, t) ** g(x, t) \quad (1.34)$$

same as above example, we make calculations and obtain

$$D(x, t) = -AC = \frac{1}{259841930424676205263257600000} x^{26} t^{40} \quad (1.35)$$

From Eq(1.35) we see that Eq(1.34) is hyperbolic for all point in $\mathbb{R}^2$.

Case (3): If $i_1 + i_2 + \dots + i_l + j_1 + j_2 + \dots + j_l$ are odd, $\alpha + \beta$ and $\mu + \nu$ are even in Eq (1.24), then we are going to classify Eq (1.24) by using Eq (1.31), in this case $b = 0$ therefore we see that the power of x, t in Eq (1.31) are even then it follows that for all point (x_0, t_0) in the domain $\mathbb{R}^2$ Eq (1.24) is Hyperbolic. As before, we consider simple non-constant wave equation in one dimensional form:

$$x^3 t^4 ** (x^6 t^5 ** x^3 t^7) u_{tt} - x^3 t^4 ** (x^6 t^5 ** x^7 t^5) u_{xx} = f(x, t) ** g(x, t) \quad (1.36)$$

and as above we see that

$$D(x, t) = \frac{1}{26409729190021827098490961920000} x^{32} t^{34} \quad (1.37)$$

Thus Eq(1.37) is positive for all points $(x_0, t_0) \in \mathbb{R}^2$; thus Eq (1.36) is hyperbolic.

Case(4): If $i_1 + i_2 + \dots + i_l + j_1 + j_2 + \dots + j_l$ are even, $\alpha + \beta$ and $\mu + \nu$ are odd in Eq (1.24), it is easy to see the power of x, t in Eq (1.31) are even, then for all point (x_0, t_0) in the domain $\mathbb{R}^2$ Eq (1.24) is Hyperbolic.

Secondly, we aim to examine the classification of generalized elliptic equation; we consider the equations with non-constant coefficients in the form of

$$a(x, y)u_{xx} + 2b(x, y)u_{xy} + c(x, y)u_{yy} = f_1(x, y) ** f_2(x, y) \quad (1.38)$$

where a, b, c and k , the polynomials and $\prod_{r=1}^z (k_r(x, y))$, the finite double convolution then Eq (1.38) becomes

$$\prod_{r=1}^z k_r(x, y) ** (a(x, y)u_{xx} + 2b(x, y)u_{xy} + c(x, y)u_{yy}) = f_1(x, y) ** f_2(x, y) \quad (1.39)$$

where $r = 1, 2, \dots, z$, and

$$\prod_{r=1}^z (k_r(x, y)) = k_1(x, y) ** k_2(x, y) ** k_3(x, y) ** \dots ** k_z(x, y)$$

where

$$k_1(x, y) = \sum_{j_1=1}^n \sum_{i_1=1}^m x^{i_1} y^{j_1}, \quad k_2(x, y) = \sum_{j_2=1}^n \sum_{i_2=1}^m x^{i_2} y^{j_2}, \quad \dots, \quad k_z(x, y) = \sum_{j_z=1}^n \sum_{i_z=1}^m x^{i_z} y^{j_z}$$

Let us compute the coefficients of Eq (1.39), by similar way that used to compute the coefficients of equation (1.24), the first coefficient given by

$$\begin{aligned} A(x, y) &= \prod_{r=1}^z k_r(x, y) * a(x, y) \\ &= \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_1=1}^n \sum_{i_1=1}^m \left(\sum_{\beta=1}^n \sum_{\alpha=1}^m P_{\alpha\beta} \frac{\Gamma\Omega}{MN} x^{i_1+i_2+\dots+i_z+\alpha+1} y^{j_1+j_2+\dots+j_z+\beta+1} \right) \end{aligned}$$

where

$$\begin{aligned} \Gamma &= i_1!i_3!\dots i_{2z-1}!, \quad \Omega = j_1!j_2!\dots j_{2z-1}!, \\ M &= ((\alpha+1)(\alpha+2)\dots(\alpha+i_1+1))((\alpha+i_1+2)(\alpha+i_1+3)\dots(\alpha+i_1+i_2+1)) \dots \\ &\quad \dots ((i_1+i_2+\dots+i_{z-1}+\alpha+2)\dots(i_1+i_2+\dots+i_z+\alpha+1)) \end{aligned}$$

and

$$\begin{aligned} N &= ((\beta+1)(\beta+2)\dots(\beta+j_1+1))((\beta+j_1+2)(\beta+j_1+3)\dots(\beta+j_1+j_2+1)) \dots \\ &\quad \dots ((j_1+j_2+\dots+j_{z-1}+\beta+2)\dots(j_1+j_2+\dots+j_z+\beta+1)) \end{aligned}$$

Similarly, the second coefficient given by

$$\begin{aligned} B(x, t) &= \prod_{r=1}^z k_r(x, y) * b(x, y) \\ &= \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_1=1}^n \sum_{i_1=1}^m \left(\sum_{t=1}^n \sum_{s=1}^m \frac{FG}{QE} x^{i_1+i_2+\dots+i_z+s+1} y^{j_1+j_2+\dots+j_z+t+1} \right) \quad (1.40) \end{aligned}$$

where

$$\begin{aligned} F &= i_1!i_3!\dots i_{2z-1}!, \quad G = j_1!j_2!\dots j_{2z-1}!t!, \\ Q &= ((s+1)(s+2)\dots(s+i_1+1))((s+i_1+2)(s+i_1+3)\dots(s+i_1+i_2+1)) \dots \\ &\quad \dots ((i_1+i_2+\dots+i_{z-1}+s+2)\dots(i_1+i_2+\dots+i_z+s+1)) \\ E &= ((t+1)(t+2)\dots(t+j_1+1))((t+j_1+2)(t+j_1+3)\dots(t+j_1+j_2+1)) \dots \\ &\quad \dots ((j_1+j_2+\dots+j_{z-1}+t+2)\dots(j_1+j_2+\dots+j_z+t+1)) \end{aligned}$$

and $s = \frac{\alpha+k}{2}$, $t = \frac{\beta+l}{2}$. Also the last coefficient given by

$$\begin{aligned} C(x, t) &= \prod_{r=1}^z k_r(x, y) * c(x, y) \\ &= \sum_{j_1=1}^n \sum_{i_1=1}^m \dots \sum_{j_1=1}^n \sum_{i_1=1}^m \left(\sum_{t=1}^n \sum_{s=1}^m q_{kl} \frac{FG}{QE} x^{i_1+i_2+\dots+i_z+k+1} y^{j_1+j_2+\dots+j_z+l+1} \right) \quad (1.41) \end{aligned}$$

where

$$\begin{aligned} F &= i_1!i_3!\dots i_{2z-1}!, \quad G = j_1!j_2!\dots j_{2z-1}!l!, \\ Q &= ((k+1)(k+2)\dots(k+i_1+1))((k+i_1+2)(k+i_1+3)\dots(k+i_1+i_2+1)) \dots \\ &\quad \dots ((i_1+i_2+\dots+i_{z-1}+k+2)\dots(i_1+i_2+\dots+i_z+k+1)) \end{aligned}$$

and

$$\begin{aligned} E &= ((l+1)(l+2)\dots(l+j_1+1))((l+j_1+2)(l+j_1+3)\dots(l+j_1+j_2+1)) \dots \\ &\quad \dots ((j_1+j_2+\dots+j_{z-1}+l+2)\dots(j_1+j_2+\dots+j_z+l+1)) \end{aligned}$$

We assume that all the coefficients $A(x, y), B(x, y), C(x, y)$ are convergent. Now we can easily check the Eq (1.39) whether it is elliptic or not. Now we have the following two cases.

(i): If $i_1 + i_2 + \dots + i_z + j_1 + j_2 + \dots + j_z, s + t, \alpha + \beta$ and $k + l$ are odd in the Eq (1.39), the power $s = \frac{\alpha+k}{2}$ and $t = \frac{\beta+l}{2}$ and the power of x and y in polynomials $a(x, y), c(x, y)$ either even or odd, also the coefficient of two polynomials $a(x, y), c(x, y)$ have the same sign, now we are going to study the classification of Eq (1.39), now the power of x, y in $B(x, y)^2 =$ the power of $A(x, y)C(x, y)$ and the coefficient of $A(x, y)C(x, y) > 1$ then the power is even, then for all point (x_0, y_0) in the domain $\mathbb{R}^2$ the Eq (1.39) is elliptic equation. For example in particular we can have

$$x^7 y^4 * [(x^3 y^2 * x^2 y^3) u_{xx} + (x^3 y^2 * x^3 y^4) u_{xy} + (x^3 y^2 * x^4 y^5) u_{yy}] = f(x, y) * g(x, y) \quad (1.42)$$

and compute the coefficient of Eq (1.42) by using Eq (1.22), Eq (1.40) and Eq (1.41), we obtain

$$D(x, y) = -\frac{23}{466987721099855155200000000} x^{30} y^{24} \quad (1.43)$$

Then it is easy to see that Eq(1.43) is always negative for all (x_0, y_0) in the domain $\mathbb{R}^2$, and thus Eq (1.42) is an elliptic equation.

(ii): If $i_1 + i_2 + \dots + i_z + j_1 + j_2 + \dots + j_z, s + t, \alpha + \beta$ and $k + l$ are even and Eq (1.39) to be elliptic equation at the condition since the power $s = \frac{\alpha+k}{2}$ and $t = \frac{\beta+l}{2}$, also the coefficients of two polynomials $a(x, y), c(x, y)$ have the same sign, thus similar to the previous section the classification of equation the Eq (1.39) on using Eq(1.22) and since the power of x and y in Eq (1.39) are even, then for all point (x_0, y_0) in the domain $\mathbb{R}^2$ Eq (1.39) is elliptic equation.

Conclusion

We note that the classification of generalized hyperbolic and elliptic equations with non-constant coefficients are still similar with original equations, that is, they are invariant after convolutions product.

REFERENCES

- [1] G. L. Lamb, Jr. Introductory Applications of Partial Differential Equations (with Emphasis on Wave Propagation and Diffusion), John Wiley and Sons, New York, 1995.
- [2] Tyn Myint-U. Partial Differential Equations of Mathematical Physics, New York, 1980.
- [3] N. Sneddon, S. Ulam and M. Stark. Operational Calculus in Two Variables and its Applications, Pergamon Press Ltd 1962.
- [4] A. Kiliçman and H. Eltayeb. A note on classification of hyperbolic and elliptic equations with polynomial coefficients, Applied Mathematics Letters 21(11)(2008), pp. 1124-1128
- [5] H. Eltayeb and A. Kiliçman. A Note on Solutions of Wave, Laplace's and Heat Equations with Convolution Terms by Using Double Laplace Transform, Applied Mathematics Letters 21(12)(2008), pp. 1324-1329.

DEPARTMENT OF MATHEMATICS AND INSTITUTE FOR MATHEMATICAL RESEARCH, UNIVERSITY PUTRA MALAYSIA, 43400 UPM, SERDANG, SELANGOR, MALAYSIA

E-mail address: akilicman@umt.edu.my, hassangadeen@hotmail.com